



Royal Netherlands
Meteorological Institute
*Ministry of Infrastructure and the
Environment*

Satellite Detection of Absorbing Aerosols

Pepijn Veefkind
Martin de Graaf
Gijsbert Tilstra
Piet Stammes

veefkind@knmi.nl


Retrieval of Aerosol Optical Depth over Land using two-angle view Satellite Radiometry during TARFOX

J. Pepijn Veefkind and Gerrit de Leeuw

TNO Physics and Electronics Laboratory, The Hague, The Netherlands

Phillip A. Durkee


Naval Postgraduate School, Monterey CA, USA



Aerosol optical depth over Europe in August 1997 derived from ATSR-2 data.



C. Robles Gonzalez, J. P. Veefkind ¹ and G. de Leeuw

TNO Physics and Electronics Laboratory, The Hague, The Netherlands




Contents

- UV Aerosol Index
- Multi-Sensor Data Record
- Absorbing aerosol over clouds
- Aerosol Index Simulator




UV Aerosol Index

- Simple quantity derived from a wavelength pair in the UV (340 – 390 nm)
- Can be used to detect plumes of absorbing aerosol
- Advantage: simple algorithm, no a priori information, very long dataset, can be applied over clouds and bright surfaces.
- Disadvantage: not a geophysical quantity which makes quantitative interpretation challenging







UV Aerosol Index Dependencies




Multi Sensor AAI Product

	Wavelength pair (nm)	Equator crossing time	Pixel size (km)	Days needed for global coverage	Platform / Operation period
TOMS	360 / 331	12 : 00 LT	50 × 50	1	Nimbus-7 (1978 – 1993)
GOME-1	340 / 380	10 : 30 LT	320 × 40	3	ERS-2 (1995 – 2003*)
SCIAMACHY	340 / 380	10 : 00 LT	60 × 30	6	Envisat (2002 – 2012)
OMI	354 / 388	13 : 45 LT	13 × 24	1	Aura (2004 – present)
GOME-2	340 / 380	09 : 30 LT	80 × 40	1.5	MetOp-A/B (2007 – present)







Differential Aerosol Absorption





Southeast Atlantic Ocean *above cloud* observations

ENVISAT track 10 Aug 2006 09:10:26 – 09:16:38 UTC



De Graaf et
al., 2012.
JGR

Southeast Atlantic Ocean *above cloud* observations





Aerosol DRE from
SCIAMACHY averaged
over the southeast
Atlantic Ocean:

- 2006 was anomalous with extreme events
- High observed values are **not reproduced** by climate models


Combining OMI and MODIS reflectances

13 Aug 2006





AAI Simulator Concept





AAI Simulator

a tool to calculate AAI from CTM model fields


- Low CPU load
- Applicable to multiple CTMs
- Applicable to multiple satellite data sets
- Accurate

GOCART




- A global atmospheric process model using assimilated meteorological fields from the Goddard Earth Observing System Data Assimilation System
- Aerosol simulations in GOCART include the major aerosol types of sea salt, sulfate, dust, black carbon, organic carbon
- The model accounts for emissions from fossil fuel and biofuel combustion, biomass burning, volcanic eruptions, vegetation, deserts, and oceans.
- Calculating aerosol composition, 4-D distributions, optical properties, radiative forcing, etc.
- See Chin et al., Atmos. Chem. Phys., 14, 3657–3690, 2014, and references therein.


OMI




SIM



OMI





GOCART + SIM




West Africa (biomass burning)

Regional mean of OMI AAI and of simulator AAI:



There is quite some agreement, even on shorter time scales. Absolute magnitude is not captured completely. The simulator underestimates the AAI values reached.

West Africa (biomass burning)





Observation Swath


■ NPP

■ S5p



TROPOMI

- ▶ UV-VIS-NIR-SWIR spectrometer.
- ▶ Spectral range: 276-775, 2305-2385 nm
- ▶ Spectral Resolution: 0.35 nm
- ▶ Spatial Resolution: 70 km
- ▶ Global daily coverage: 1 day



Copernicus

CO₂, CH₄, HCHO
column

Clouds & Aerosol layer height

Summary

- The UV spectrometers can provide information on absorbing aerosol plumes.
- The UV AI has strong dependency on the absorption, the concentration and height of aerosols.
- We can use the UV observations in a quantitative way:
 - For biomass burning plumes over clouds
 - Using simulators
- At KNMI we will extend the UV AI with the European UVN spectrometers: TROPOMI, Sentinel 4 and Sentinel 5
 - For these instrument we are also developing aerosol layer height retrievals